

# **SYNTHESIS, CHARACTERIZATION AND BIOLOGICAL EVALUATION OF SOME NEWER SERIES OF QUINAZOLONES**

Dissertation submitted to

The Tamil Nadu Dr. M.G.R. Medical University,

Chennai- 600032.

In the partial fulfillment for the award of Degree of

**MASTER OF PHARMACY**

(Pharmaceutical Chemistry)

Submitted by

**SANKARNARAYANAN. S**

Reg.No. 26106035

Under the Guidance of

**Mr. A. THIRUGNANA SAMBANTHAN, M.Pharm., (Ph.D.)**

Assistant Professor,

Department of Pharmaceutical Chemistry,



**ADHIPARASAKTHI COLLEGE OF PHARMACY,**

(Accredited By “NAAC” with CGPA of 2.74 on a Four point Scale at “B” Grade)

**MELMARUVATHUR- 603319.**

**MAY- 2012.**

# **SYNTHESIS, CHARACTERIZATION AND BIOLOGICAL EVALUATION OF SOME NEWER SERIES OF QUINAZOLONES**

Dissertation submitted to

The Tamil Nadu Dr. M.G.R. Medical University,

Chennai- 600032.

In the partial fulfillment for the award of Degree of

**MASTER OF PHARMACY**

(Pharmaceutical Chemistry)

Submitted by

**SANKARNARAYANAN. S**

Reg.No. 26106035

Under the Guidance of

**Mr. A. THIRUGNANA SAMBANTHAN, M.Pharm., (Ph.D.)**

Assistant Professor,

Department of Pharmaceutical Chemistry,



**ADHIPARASAKTHI COLLEGE OF PHARMACY,**

(Accredited By “NAAC” with CGPA of 2.74 on a Four point Scale at “B” Grade)

**MELMARUVATHUR- 603319.**

**MAY- 2012.**

## **CERTIFICATE**

This is to certify that the dissertation entitled **“SYNTHESIS, CHARACTERIZATION AND BIOLOGICAL EVALUATION OF SOME NEWER SERIES OF QUINAZOLONES”** submitted to The Tamil Nadu Dr. M.G.R. Medical University in partial fulfillment for the award of the Degree of Master of Pharmacy (Pharmaceutical Chemistry) was carried out by **SANKARNARAYANAN. S** (Reg. No. 26106035) in the Department of Pharmaceutical Chemistry under my direct guidance and supervision during the academic year 2011-2012.

Place: Melmaruvathur

Date:

**Mr. A. THIRUGNANA SAMBANTHAN, M.Pharm., (Ph.D.)**

Assistant Professor,

Department of Pharmaceutical Chemistry,

Adhiparasakthi College of Pharmacy,

Melmaruvathur- 603319.

## **CERTIFICATE**

This is to certify that the dissertation entitled “**SYNTHESIS, CHARACTERIZATION AND BIOLOGICAL EVALUATION OF SOME NEWER SERIES OF QUINAZOLONES**” is the bonafide research work carried out by **SANKARNARAYANAN. S** in the Department of Pharmaceutical Chemistry, Adhiparasakthi College of Pharmacy, Melmaruvathur which is affiliated to The Tamilnadu Dr. M.G.R Medical University under the guidance of **Mr. A. THIRUGNANA SAMBANTHAN, M.Pharm., (Ph.D.)** Assistant Professor, Department of Pharmaceutical Chemistry, Adhiparasakthi College of Pharmacy, during the academic year 2011-2012.

Place: Melmaruvathur

Date:

**Prof. (Dr.). T. VETRICHELVAN, M.Pharm., Ph.D.,**  
Principal,  
Adhiparasakthi College of Pharmacy,  
Melmaruvathur- 603319.

## ACKNOWLEDGEMENT

First and foremost, I wish to express my deep sense of gratitude to **HIS HOLINESS ARULTHIRU ADIGALAR**, President, ACMEC Trust, Melmaruvathur for his ever growing blessings in each step of the study.

Words are not just enough to express our gratitude to the blessing of **GOD** throughout the work. Indeed our project is a small work done with the help of primitive persons at heart. So it is my bounded duty to promulgate them individually.

With great respect and honor, I extend my thanks to **Thirumathi LAKSHMI BANGARU ADIGALAR**, Vice President, ACMEC Trust, Melmaruvathur for this opportunity and support all the way in completing the study.

I got inward bound and brainwave to endure experimental investigations in synthetic chemistry to this extent; I concede my most special gratitude and thanks to **Mr. A. THIRUGNANA SAMBANTHAN, M.Pharm., (Ph.D.)** Assistant Professor, Department of Pharmaceutical Chemistry, Adhiparasakthi College of Pharmacy for the active guidance, valuable suggestions and a source of inspiration for my work.

I owe my sincere thanks with bounteous pleasure to **Prof. (Dr.) T. VETRICHELVAN, M.Pharm., Ph.D.**, Principal, Adhiparasakthi College of Pharmacy, without his encouragement and supervision it would have been absolutely impossible to bring out the work in this manner.

I have great pleasure in expressing my sincere heartfelt thanks to **Prof (Dr.) D. Nagavalli, M.Pharm., Ph.D.** and **Mr. M. Sugumaran, M.Pharm., (Ph.D.)** Associate Professor Department of Pharmaceutical Chemistry for their encouragement and support for the successful completion of this work.

My sincere thanks to our lab technicians **Mrs. S. Karpagavalli, D.Pharm., Mr. M. Gomathi Shankar, D.Pharm., Mrs. N. Thatchayani, D.Pharm.,** and office assistant **Mr. I. Kumar** for their kind help throughout this work.

I extend my great thankful to **Mrs S. Shoba M.Pharm., (Ph.D.)** Assistant professor, **Ms. M. Vijayakumari M.Pharm., Lecturer,** Department of Pharmacology, for their help during analgesic activity study.

I am indeed very much thankful to the librarian **Mr. M. Suresh, M.L.I.S.,** for providing all reference books for completion of this project.

I am thankful to all my colleagues, juniors, seniors and my friends for their support and suggestion during my work.

Finally yet importantly, I one word forward my affectionate thanks to my family members, especially my parents, brother and sisters for their frequent prayers which has sustained me a lot in the successful completion of my project work.

**SANKARNARAYANAN. S**

DEDICATED TO MY  
PARENTS AND GOD

## CONTENTS

S. No	Content	Page. No
I	INTRODUCTION	1
II	OBJECTIVES OF THE WORK	7
III	LITERATURE REVIEW	8
IV	EXPERIMENTAL SECTION	
A	Materials and methods	22
B	Scheme of the work	25
C	Methodology	27
D	Evaluation of Analgesic Activity	31
E	Evaluation of Antibacterial Activity	33
F	Evaluation of Antifungal Activity	35
V	RESULTS AND DISCUSSION	37
VI	SUMMARY	123
VII	CONCLUSION	125
VIII	BIBLIOGRAPHY	126
IX	ANNEXURE-I	132
	ANNEXURE-II	133



## LIST OF TABLES

Table No	Content	Page No
1	List of Chemicals and Instruments Used	22
2	Physical Data of BOC Amino acids	37
3	Physical Data of 2-chloromethyl-3-amino acid substituted quinazolin-4-(3H)-ones	38
4	Physical Data of 2-chloromethyl-3-heterocyclic substituted quinazolin-4(3H)-ones	38
5	Interpretation of IR spectra of QAL	39
6	Interpretation of IR spectra of QAA	41
7	Interpretation of IR spectra of QAPA	43
8	Interpretation of IR Spectra of QAG	45
9	Interpretation of IR Spectra of QBEN	47
10	Interpretation of IR Spectra of QIBM	49
11	Interpretation of IR Spectra of QNIBM	51
12	Interpretation of IR Spectra of QPIN	53
13	Interpretation of IR Spectra of QBrP	55
14	Interpretation of NMR Spectra of QAL	57
15	Interpretation of NMR Spectra of QAA	63
16	Interpretation of NMR Spectra of QAPA	67
17	Interpretation of NMR Spectra of QAG	73
18	Interpretation of NMR Spectra of QBEN	77
19	Interpretation of NMR Spectra of QIBM	81
20	Interpretation of NMR Spectra of QNIBM	84
21	Interpretation of NMR Spectra of QPIN	89
22	Interpretation of NMR Spectra of QBrP	92
23	Interpretation of Mass Spectra of QAL	97
24	Interpretation of Mass Spectra of QAA	99
25	Interpretation of Mass Spectra of QAPA	101
26	Interpretation of Mass Spectra of QAG	103
27	Interpretation of Mass Spectra of QBEN	105
28	Interpretation of Mass Spectra of QIBM	107
29	Interpretation of Mass Spectra of QNIBM	109

30	Interpretation of Mass Spectra of QPIN	111
31	Interpretation of Mass Spectra of QBrP	113
32	Results of Analgesic activity	115
33	Results of Antibacterial activity	116
34	Results of Antifungal activity	119

## LIST OF FIGURES

Figure No	Content	Page No
1	Scheme I	25
2	Scheme II	26
3	I.R Spectra of QAL	40
4	I.R Spectra of QAA	42
5	I.R Spectra of QAPA	44
6	I.R Spectra of QAG	46
7	I.R Spectra of QBEN	48
8	I.R Spectra of QIBM	50
9	I.R Spectra of QNIBM	52
10	I.R Spectra of QPIN	54
11	I.R Spectra of QBrP	56
12	NMR Spectra of QAL	58
13	NMR Spectra of QAL	59
14	NMR Spectra of QAL	60
15	NMR Spectra of QAL	61
16	NMR Spectra of QAL	62
17	NMR Spectra of QAA	64
18	NMR Spectra of QAA	65
19	NMR Spectra of QAA	66

20	NMR Spectra of QAPA	68
21	NMR Spectra of QAPA	69
22	NMR Spectra of QAPA	70
23	NMR Spectra of QAPA	71
24	NMR Spectra of QAPA	72
25	NMR Spectra of QAG	74
26	NMR Spectra of QAG	75
27	NMR Spectra of QAG	76
28	NMR Spectra of QBEN	78
29	NMR Spectra of QBEN	79
30	NMR Spectra of QBEN	80
31	NMR Spectra of QIBM	82
32	NMR Spectra of QIBM	83
33	NMR Spectra of QNIBM	85
34	NMR Spectra of QNIBM	86
35	NMR Spectra of QNIBM	87
36	NMR Spectra of QNIBM	88
37	NMR Spectra of QPIN	90
38	NMR Spectra of QPIN	91
39	NMR Spectra of QPIN	93
40	NMR Spectra of QBrP	94
41	NMR Spectra of QBrP	95
42	NMR Spectra of QBrP	96
43	Mass Spectra of QAL	98
44	Mass Spectra of QAA	100
45	Mass Spectra of QAPA	102

46	Mass Spectra of QAG	104
47	Mass Spectra of QBEN	106
48	Mass Spectra of QIBM	108
49	Mass Spectra of QNIBM	110
50	Mass Spectra of QPIN	112
51	Mass Spectra of QBrP	114
52	Antibacterial activity against <i>S.aureus</i>	117
53	Antibacterial activity against <i>E.coli.</i>	118
54	Antifungal activity against <i>A.niger</i>	120

## ABBREVIATIONS

Ala	-	Alanine
Gly	-	Glycine
Phe	-	Phenyl alanine
Leu	-	Leucine
BOC	-	Tert-butyloxy carbonic anhydride
IPA	-	Iso propyl alcohol
DCM	-	Dichloro methane
THF	-	Tetrahydro furan
TEA	-	Triethyl amine
EDC	-	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride
RT	-	Room temperature
M.P	-	Melting point
R <sub>f</sub>	-	Retardation factor
m.mol	-	Millimole
g	-	Gram
ml	-	Milli litre
ppm	-	Parts per million
IR	-	Infra Red spectroscopy
<sup>1</sup> HNMR	-	Proton Nuclear Magnetic Resonance spectroscopy
<sup>0</sup> C	-	Degree Celsius
h	-	Hour
min	-	Minutes

# INTRODUCTION

## I. INTRODUCTION

The name quinazoline (German: Chinazolin) was first proposed for this compound by Weddige, on observing that this was isomeric with the compounds cinnoline and quinoxaline. Paal and Bush suggested the numbering of quinazoline ring system, which is currently used. The other less commonly used names for this ring system are 'phenmiazine' and 5,6-benzopyrimidine.

Of many derivatives of quinazoline system known so far, keto-quinazolines also called as quinazolinones, are the most important compounds. Depending upon the position of the keto or oxo group, these compounds may be classified into two types: 2-(1H) quinazolinones (or) 1, 2-dihydro-2-oxo quinazolines and 4(3H)-quinazolines or 3, 4-dihydro-oxoquinazolines. These systems exhibit lactam-lactim tautomerism and undergo hydroxyl group replacement reactions. 2-Cyano-4(3H)-quinazolinone was the first quinazolinone derivative to be synthesized.

### Reactivity of 4(3H)-Quinazolinones

Reactions associated with tautomeric nature of the quinazolinones are often quite complex and generally unpredictable. The recorded chemical investigation on the subject is voluminous. The amide linkages in quinazolinones should not be looked on as predominantly the keto or the enol form but as true keto-enol tautomers, showing reaction characteristic of both the forms.

Quinazolinones are always high melting crystalline solids, insoluble in water and in most organic solvents but soluble in aqueous alkali. They are generally insoluble in dilute acids but are sometimes soluble in concentrated acids. Simple 4(3H)-quinazolinones, although insoluble in dilute acids, are soluble in 6N hydrochloric acid. 4(3H)-quinazolinones form stable monohydrochlorides,

chloroplatinate, chloroaurates and picrates and their metal salts of silver, mercury, zinc, copper, sodium and potassium.

### **Stability of the ring system**

The ring system in quinazolinone is exceedingly stable in oxidation, reduction, hydrolysis reactions and other treatment designed to break the ring. There is no report of degradation of quinazolinone by simple chemical oxidation.

### **Biological importance of 4(3H)-Quinazolinones**

The quinazolinone skeleton is a frequently encountered heterocycle in medicinal chemistry literature with applications including antibacterial, analgesic, antiinflammatory, antifungal, antimalarial, antihypertensive, CNS depressant, anticonvulsant, antihistaminic, local anaesthetic, antiparkinsonism, antiviral and anticancer activities. Little number of quinazolinones was reported as potent chemotherapeutic agents in the treatment of tuberculosis. For example 3-aryl-6, 8-dichloro-2*H*-1,3-benzoxazin-2,4(3*H*)-diones and 3-arylquinazolin-2,4(1*H*,3*H*)-diones as antimycobacterial agents, quinazolinone derivatives as antitubercular agents. (Dr. S. N. Meyyanathan, *Pharmainfo.net.*, 2005).

Research programs for the discovery of new antimicrobial drug for improving the evaluation criteria are under way in many laboratories. In addition, knowledge of specific constituents of the mycobacterial cell and their biochemical role has advances considerably in the recent years and may permit a more rational approach to the design of new drug acting on specific targets. Also recent improvements in the knowledge of the mechanism of action of the available drug in the biochemical mechanism of resistance to them may be used as a basis for designing new and better weapons to fight the mycobacterial diseases.



The quinazolines and quinazolones when selectively functionalized act as building blocks for the preparation of numerous biological active compounds. The quinazolines are found to exhibit antifungal, antitubercular, antitumour and anticonvulsant activities. (*Shahikant R Pattan., et al., 2006*).

The need of new antimicrobial agents is justified because more microorganisms are being resistance to the present drugs available in the market. Worldwide researchers are trying to synthesize new drugs with better pharmacokinetic and dynamic properties with less adverse effects. The literature survey suggests that the 2, 3-substituted quinazolin-4-ones have proved to be the good bioactive molecules. They have shown diverse biological activities like antibacterial, antifungal, antiinflammatory, antitubercular, anticonvulsant, antiHIV, cardiac stimulant, diuretic and anticancer activity etc. (*Jayshari S Pattan., et al., 2009*).

Pharmacologically, quinazolin-4-ones are most important classes of heterocyclic compounds. These compounds possess versatile type of biological activities. Structure activity relationship studies of quinazolinone ring system revealed in various literatures suggest that position 2, 6 and 8 are very much important for structure activity studies. (*Raghavendra N.M., et al., 2008*).

The exploitation of a simple molecule with different functionalities for the synthesis of heterocyclic is a worthwhile contribution in the chemistry of heterocycles. Quinazolones have been frequently used in medicine because of their wide spectrum of biological activities. In light of the growing number of applications in recent years, there has been an enormous increase in the interest among biologists and chemists in their synthesis and bioactivity of quinazoline derivatives.

Analgesics are the drug which relieves pain without disturbing consciousness. Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage. Pain acts as a warning signal against disturbances either in the body or in the external environment of an individual and thus has protective function. (*Gupta S. K ., 2004*)

Analgesics are classified into opioid and Non opioid analgesics. The term opioid is used to denote all naturally occurring, semisynthetic and synthetic drugs, which have morphine like action. These drugs are called Narcotic analgesics. e.g. morphine and their derivatives, codeine and its derivatives, synthetic compounds such as Pethidine, Methadone, Propoxyphene, Levorphanol and Tramadol.

Opioid drugs produce their effects by binding to opioid receptors, which are widely distributed in CNS and other tissues. Opioid receptors are also present in the peripheral nerves where they probably respond to peripherally applied opioids and locally released endogenous peptides during inflammation.

Non opioid analgesics are the agents which do not interact with opioid receptors and relieve pain without depressing the CNS. e.g. Salicylates and related compounds. Efforts towards the development and identification of new molecules for analgesic and antiinflammatory activities with minimal gastrointestinal ulceration side effects have gained lot of significance. There are no promising quinazolines which are in the market in these NSAIDS category except few drugs like Proquazone, Afloqualone and Diproqualone etc.

The novel derivatives of quinazolines mentioned might be beneficial in terms of biological activity for which further studies can be done to confirm it as a potential drug candidate. Proquazone a derivative of quinazolin-4-one exhibits potential NSAID which has been used in the disease conditions like rheumatoid arthritis,

ankylosing spondylitis, osteoarthritis, inflammatory conditions and acute pain states such as dysmenorrhea, postoperative pain and headache. Its derivative fluproquazonal has potent analgesic action. (Vijaychand *et al.*, 2011)

Bacteria are microorganisms that have circular double-stranded DNA and (except for mycoplasmas) cell walls. Most bacteria live extracellularly. Some bacteria eg. *Salmonella typhi*, *Neisseria gonorrhoeae*, *Legionella*, *Mycobacterium*, *Chlamydia*, and *Chlamydophila* spp preferentially reside and replicate intracellularly. Some bacteria such as chlamydiae and rickettsiae are obligate intracellular pathogens (i.e., able to grow, reproduce, and cause disease only within the cells of the host). Others (eg, *Salmonella typhi*, *Brucella* sp, *Francisella tularensis*, *N. gonorrhoeae*, *N. meningitidis*, *Legionella* and *Listeria* spp, *Mycobacterium tuberculosis*) are facultative intracellular pathogens.

Antibacterial drugs are derived from bacteria or molds or are synthesized by de novo. Technically, antibiotic refers only to antimicrobials derived from bacteria or molds but is often including used synonymously with antibacterial drug.

Antibiotics have many mechanisms of action, including inhibiting cell wall synthesis, activating enzymes that destroy the cell wall, increasing cell membrane permeability and interfering with protein synthesis and nucleic acid metabolism.

Antibiotics sometimes interact with other drugs. Raising or lowering serum levels of other drugs by increasing or decreasing their metabolism or by various other mechanisms.

The rapid rise in bacterial resistance to the traditional antibiotics such as penicillins and tetracyclines has encouraged a continuing search for new classes of

compounds with novel modes of antibacterial activity.

In view of the above reports, the design and synthesis of newer antimicrobials is an area of immense significance and continues to attract the attention of increasing number of medicinal chemists.

With the interest of on quinazolin-4-ones due to its extension pharmacological actions we have designed a new series of quinazolinone substituted at 2<sup>nd</sup> position with chloromethyl and 3<sup>rd</sup> position with different amino acid and heterocycles.

# OBJECTIVE OF THE WORK



## **II. OBJECTIVES OF THE WORK**

- To synthesize 2-chloromethyl-3-amino acid substituted quinazolin-4(3*H*)-one.
- To synthesize different 2-chloromethyl-3-heterocyclic substituted Quinazolones by cyclizing the chloroacetyl anthranilic acid with amino substituted indole, benzthiazole and pyridine.
- To characterize the structures of the synthesized compounds by various spectroscopic methods via IR, NMR and Mass spectroscopy.
- To screen the analgesic activity of synthesized compounds by Eddy's hot plate method.
- To evaluate the antimicrobial activity of synthesized compounds by Disc Diffusion method.
- To interpret and conclude the results obtained from the study.




# LITERATURE REVIEW

### III. LITERATURE REVIEW

**Rajitha G. *et al.*, (2011)** synthesized some novel 3-[5-phenyl-1,3,4-oxadiazol-2-yl]-2-(substituted styryl)-quinazolin-4(3H)-ones and evaluated for antibacterial activity.




**Narendra babu *et al.*, (2011)** carried out the synthesis of new quinazolinone and the compounds were screened for antiinflammatory and anthelmintic activities.






**Ponnilarasan Ilangoan *et al.*, (2010)** synthesized a series of different quinazolinone derivatives, and they were examined by invitro studies for anticonvulsant and antimicrobial activities.




**Mariappan G. *et al.*, (2010)** synthesized a series of some unique quinazolone fused schiff bases and those compounds were evaluated for biological activity.







**Rajveer C. H. *et al.*, (2010)** carried out the synthesis of 2-(6-bromo-2-phenyl-4-oxoquinazolin-3(4H)-yl)-N-substituted acetamides and 1-amino-5-(6-bromo-3, 4-dihydro-2-phenyl-4-oxoquinazolin-3yl) methyl-1,3,4-triazin-2-thiol and the selected compounds were examined for antimicrobial activity.




**Dubey *et al.*, (2010)** synthesized a novel series of 2-(4-substituted sulfonyl piperazin-1-yl)methyl-3-aryl-quinazolin-4(3H)-one and screened for the antimicrobial activity.



**Shashikant R. Pattan *et al.*, (2009)** synthesized some new quinazolinone derivatives and the proposed compounds were screened for antimicrobial, antifungal, antitubercular and antiinflammatory activities.



**Narsaiah B. *et al.*, (2009)** carried out the synthesis of novel 2, 4, 6 trisubstituted quinazoline derivatives showing antibacterial and cytotoxic activities against THP-1, HL-60 and A375 cell line.





$\text{X} = \text{H}, \text{Br}, \text{I}$




$\text{R} = \text{CH}_3, \text{CF}_3, \text{C}_6\text{H}_5, \text{C}_6\text{H}_4\text{F}_3$

**Vijayanand R. *et al.*, (2009)** synthesized a series of 4-oxo-2-phenyl-4H quinazolin-3-carboxylic acid (4-substituted phenyl amides) by condensing 2-phenyl-3, 1-


benzoxazin-4one and 4-substituted phenyl urea. The compounds were found to have significant effect against bacterial and fungal organisms.





**Salahuddin M. D. *et al.*, (2009)** synthesized some novel 3-(6-Substituted-1, 3-benzothiazol-2-yl)-2-[(4-substituted phenyl) amino] methyl quinazolin-4(3H)-ones with various substituted amine and selected derivatives are investigated for antiinflammatory and antibacterial activities.




**Gowramma B. *et al.*, (2009)** synthesized few compounds of 1-substituted phenyl 2-[(bis-(2-chloroethyl)amino)-methyl-7-nitro-4(1H)-quinazolinone by chlorination with phosphorous oxychloride and phosphorous penta chloride and all the compounds were incorporated with nitrogen mustard moiety and screened for anticancer activity.



**Avinsah patil *et al.*, (2009)** carried out a new synthesis of 2-[[5-substituted-1H-benzimidazol-2-yl sulfinyl] methyl]-3-substituted Phenyl Quinazolin-4(3H)-one derivatives; antiulcer activity of the derivatives were studied by pylorous ligation induced ulcer models in rats.




**Jai singh *et al.*, (2008)** synthesized an innovative series of 2-{5'-(4-Pyridinyl)-1',2',3'-oxadiazol-2-yl-thiomethyl}-3-substituted-aryl-6-substituted-quinazolin-4-one and all the compounds were screened for antiinflammatory activity.



X = H, Br, I


R = H, Cl, OCH<sub>3</sub>,

**Raghavendra M. N. *et al.*, (2008)** synthesized by condensation of 2-chloro-*N*-(4-oxo-substituted-quinazolin-3(4*H*)-yl)-acetamides with various substituted piperazines through single step reaction and the derivatives were investigated for antibacterial and antifungal activities.




X = H, Br

R = CH<sub>2</sub>, CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>3</sub>,



**Suthakaran R. *et al.*, (2008)** synthesized a different sequence of 3-(2-(4)-4-substituted benzylidene-4, 5-dihydro-5-oxo-2-phenyl imidazol-1-yl) ethyl)-6, 8-un/dibromo substituted-2-substituted quinazolin-(3*H*)-one and compounds were investigated for antibacterial and antifungal activities





X = H, Br

R = CH<sub>3</sub>, CH<sub>2</sub>Cl, C<sub>6</sub>H<sub>5</sub>


R<sub>1</sub> = OH, Cl

**Srinivasa reddy *et al.*, (2007)** synthesized a series of 1-phenyl-2-(substituted)-4-(1H) quinazolinones and the compounds were screened for in vivo analgesic and antiinflammatory activities.





**Jessy E. M. *et al.*, (2007)** synthesized a unique series of 2, 3-disubstituted-3, 1-quinazolin-4-(3H) ones and the compounds were screened for antibacterial and antiinflammatory activities.




$R = R_1 = H, COOH$

$R_2 = H, Br$

$R_3 = R_4 = H, Cl$


$R_5 = NO_2, OCH_3, OH$

**Ashis Kumar Nanda *et al.*, (2007)** synthesized of ten 3-(arylidene amino)-2-phenylquinazolin-4(3H)-ones with QSAR studies are carried out and the compounds were investigated for antimicrobial activity.





$R = CH_3, OH, F, Cl, NO_2.$

**Shashikant R. Pattan *et al.*, (2006)** synthesized a new series of N-3[(4-(4-chlorophenyl)thiazole-2-yl)-(2-(amino) methyl)-quinazolin-4(3H)-ones and all the derivatives were screened for their antitubercular activity.




**Tripti singh *et al.*, (2006)** synthesized a series of heterocyclic derivatives of quinazolinone and the compounds were screened for insecticidal and antimicrobial activities.









**Mistry D. *et al.*, (2006)** synthesized of newer quinazolinones and the compounds were screened for antimicrobial activity.






R = H, OCH<sub>3</sub>, Br, Cl, NO<sub>2</sub>,

**Manish Patel *et al.*, (2005)** carried out the synthesis of new 2, 3, 6-trisubstituted quinazolin-4(3H)-ones and antibacterial potential was evaluated for the selective moieties.










**Pandey D. *et al.*, (2005)** synthesized a new series of isoquinolinyl quinazolones and investigated all the compounds for antiviral and antifungal activities.







**Ashok kumar *et al.*, (2003)** synthesized some new 2, 3, 6 trisubstituted quinazolinone for potent antiinflammatory, analgesic and COX-2 inhibitory activities.



**Bhatta V. *et al.*, (1998)** synthesized new sequences of 3-aryl-2-(4'-aryl thiazol-2'-yl aminomethyl quinazolin-4(3H)-ones and all the compounds were investigated for antimicrobial activity.



**Surendara bahadur *et al.*, (1982)** synthesized of some newer-4(3H)-quinazolinones and the compounds were examined for antimicrobial agents.



## IV. EXPERIMENTAL SECTION

### A. MATERIALS AND METHODS

List of chemicals and instruments used for the study:

**Table-1**

S. No	Chemicals / Instruments	Supplier / Model
<b>Chemicals</b>		
1.	Alanine	Spectrochem Pvt. Ltd.,
2.	Glycine	Lobachem Pvt. Ltd.,
3.	Leucine	Spectrochem Pvt. Ltd.,
4.	Phenyl alanine	Lobachem Pvt. Ltd.,
5.	Anthranilic acid	Sigma Aldrich
6.	Chloroacetyl chloride	Lobachem Pvt. Ltd.,
7.	Benzene	Lobachem Pvt. Ltd.,
8.	Hydrazine hydrate	Lobachem Pvt. Ltd.,
9.	Ethanol	Lobachem Pvt. Ltd.,
10.	Acetone	Lobachem Pvt. Ltd.,
11.	Pyridine	Lobachem Pvt. Ltd.,
12.	Benzimidazole	Lobachem Pvt. Ltd.,
13.	Glacial acetic acid	Spectrochem Pvt. Ltd.,
14.	t-butyl oxy carbonic anhydride	Qualigens Fine Chemicals
15.	Iso propyl alcohol	Lobachem Pvt. Ltd.,
16.	Chloroform	Spectrochem Pvt. Ltd.,
17.	Petroleum ether (40-60 <sup>0</sup> c)	Qualigens Fine Chemicals

18.	Thionyl chloride	Spectrochem Pvt. Ltd.,
19.	Sodium hydroxide	Spectrochem Pvt. Ltd.,
20.	Methanol	Qualigens fine Chemicals
21.	Dichloro methane	Lobachem Pvt. Ltd.,
22.	Tetrahydro furan	Lobachem Pvt. Ltd.,
23.	Trifluoroacetic acid	Qualigens Fine Chemicals
24.	Sodium sulphate anhydrous	Qualigens Fine Chemicals
25.	Conc. sulphuric acid	Lobachem Pvt. Ltd.,
26.	Conc. hydrochloric acid	Qualigens Chemicals
27.	Diethyl ether	Qualigens Fine Chemicals
<b>Instruments</b>		
28.	Magnetic stirrer	Remi Equipments
29.	Weighing balance	Shzimadzhu 220 v
30.	Melting point apparatus	Sun bim Equipments
31.	Hot air oven	Picses Instruments
32.	Heating mantle	Ajay Equipments

## Methods

- Melting points were determined by using open capillary tubes and were uncorrected.
- The completion of reaction and purity of the compounds were checked by thin layer chromatography on pre-coated silica gel plates using iodine vapours as detecting agent.

- The I.R spectra of the synthesized compounds were recorded in JASCO FT-IR spectrophotometer in Ideal Analytical Lab, Puducherry.
- The NMR spectra of the synthesized compounds were recorded in BRUKER 400 MHz NMR spectrometer, SAIF, Punjab University, Punjab.
- The Mass spectra of the synthesized compounds are recorded in JEOL GCmate using Electron impact method as ionization mode in SAIF, IIT-Madras.
- The analgesic activity of the synthesized compounds was screened by Eddy's Hot plate method in Adhiparasakthi College of Pharmacy, Melmaruvathur.
- The antimicrobial activity of the synthesized compounds was screened by Pharma Analytical Lab, Puducherry.



Figure-1

## B. SCHEME OF THE WORK

SCHEME I





Figure-2

SCHEME II




### C. METHADODOLOGY

#### Procedure for scheme I


##### Synthesis of N-chloroacetyl anthranilic acid:

0.1M (13.7gm) of anthranilic acid was taken in a 250 ml round bottom flask followed by addition of 0.15M (16.95ml) chloroacetyl chloride in 80ml of benzene and two to three drops of pyridine under cold condition. The reaction mixture is refluxed for 6 h cooled and filtered it. The solid was purified by recrystallization from acetone-ethanol mixture (1:1). ( Shashikant R Pattan., et al., 2006)



##### Synthesis of 3-amino-2-(chloromethyl) quinazolin-4(3H)-one:

N-Chloroacetyl anthranilic acid (0.01M) was refluxed with Hydrazine hydrate (0.01M) in the presence of 10g of  $\text{K}_2\text{CO}_3$  in 100ml of dry ethanol for 16 h. The ethanol was evaporated and the mixture was neutralized with 0.1M HCl under ice cold condition. Then the residue was washed thoroughly with boiling water and recrystallized from acetone: ethanol mixture (1:1). ( Shashikant R Pattan., et al., 2006)




N-Chloroacetyl anthranilic acid Hydrazine hydrate 3-amino-2-(chloromethyl) quinazolin-4(3H)-one)

### Protection of Amino Group

The amino acid (20 m.mol) was dissolved in a mixture of 1N NaOH (20 ml) and IPA (20 ml). To this BOC (26 m.mol, 6 ml) in IPA (10 ml) was added, followed by 1N NaOH (20 ml) to the resulting solution. The solution was stirred at room temperature for 2 h, washed with light petroleum ether (40 – 60°C) (20 ml), acidified to pH-3 with 2N sulphuric acid and finally extracted with chloroform (3 x 20 ml). The organic layer was dried over anhydrous sodium sulphate and evaporated under pressure to give the BOC-amino acid. The crude product was recrystallized using a mixture of chloroform and petroleum ether. (*Bodanszky. M., 1984*)

### Coupling of quinazalone with BOC-amino acid

3-amino-2-(chloromethyl) quinazolin-4(3*H*)-one (10 m.mol) was dissolved in DCM (20 ml). To this triethyl amine (4 ml) was added at 0°C and the reaction mixture was stirred for 15 minutes. BOC- amino acid (10 m.mol) in chloroform (20 ml) and EDC (10 m.mol) were added with stirring. After 8 h the reaction mixture was filtered and residue was washed with chloroform (30 ml) and added to the filtrate. The filtrate was washed with 5% sodium bicarbonate (20 ml) and saturated sodium chloride (20 ml) and water. Organic layer was dried over anhydrous sodium sulphate and the product was recrystallized by chloroform and petroleum ether. (*Bodanszky. M., 1984*)




3-amino-2-(chloromethyl) quinazolin-4(3*H*)-one


**Synthesis of 3-amino acid substituted-2-chloromethyl quinazolin-4(3H)-one**

The protected amino acid was dissolved in chloroform (15 ml) and treated with trifluoroacetic acid (2 m.mol). The solution was stirred at room temperature for 1h and washed with saturated sodium bicarbonate (5 ml) and the organic layer was dried over anhydrous sodium sulphate and concentrated under reduced pressure.

(Bodanszky, M., 1984)


**Procedure for scheme II****Synthesis of (1H-benzimidazol-2-yl) aniline:**

A mixture of O-phenylene diamine 3.8 g (34 mM) and 4-amino benzoic acid 4.5 g (33 mM) were stirred in a syrupy ortho-phosphoric acid (45 ml) at 200° C for 2 h. The reaction mixture was cooled and poured on to the crushed ice. The bulky white precipitate obtained was stirred in cold water (400ml) and sodium hydroxide solution (5 M) was added until the pH 7. The resulting solid was filtered and recrystallized from methanol. (Sreena *et al.*, 2009)




**Synthesis of 4-(5-nitro-1H-benzimidazol-2-yl) aniline:**

Concentrated nitric acid (7.5 ml) was placed in 3-necked round bottom flask fitted with a mechanical stirrer. The flask was immersed in ice cold water and added slowly concentrated sulphuric acid (7.5 ml) down the condenser with slow stirring. After the addition, (1H-benzimidazol-2-yl) aniline 5.85 g (0.028 M) was added in a portion over a period of 1 h at such a rate that the temperature did not exceed 35° C. After continuous stirring for 12 h, the reaction mixture was poured very slowly over crushed ice with vigorous stirring. The formed product was filtered, washed with cold water and recrystallized from ethanol. (*Jitender Singh et al., 2010*)

**2-chloromethyl-3- substituted quinazolin-4(3H)-one:**

N-chloroacetyl anthranilic acid (0.01M) was refluxed with (0.01M) of 2-amino benzthiazole, 2- amino benzimidazole, 2-amino 5-nitro phenyl benzimidazole, 2-amino 5-nitro pyridine and 2-amino 5-bromo pyridine in the presence of 10g of  $\text{K}_2\text{CO}_3$  in 100ml of dry ethanol. The ethanol was evaporated and the mixture was neutralized with 0.1M HCl under ice cold condition. Then the residue was washed thoroughly with boiling water and recrystallized from acetone: ethanol mixture (1:1).



## D. EVALUATION OF ANALGESIC ACTIVITY

The drug is reflected to obligate opioid analgesic activity if it relieves pain by acting centrally to elevate pain threshold without disturbing consciousness. If it is peripherally acting analgesics, it inhibits prostaglandin synthesis and raises the threshold to pain perception. Numerous methods for evaluation of analgesic activity are available. They can be categorized into chemical, electrical, mechanical and thermal methods. (*Gupta S.K., 2004*)

Chemical Method: Writhing Test, Randall - Selitto Test and Intra Arterial Bradykinin Test.

Electrical Method: Electrical Stimulation of the Tail, Flinch Jump test in Mice and Tooth Pulp Stimulation in Rabbits


Mechanical Method: Haffner's Tail-clip test in Mice

Thermal Method: Radiant Heat Method, Tail-Immersion Test and Hot plate Method.

The thermal method was followed to screen the analgesic activity. In this method the pain is induced by heat from Eddy's Hot plate.

### Standard Drug for Analgesic Activity

To screen and compare the analgesic activity of synthesized compounds, the pentazocine was used as a standard drug.

**Standard drug**

Pentazocine

Pentazocine is the one of the potent opioid analgesic. It relieves moderate to severe pain by acting centrally to elevate pain threshold without disturbing consciousness.

**Procedure for Eddy's Hot Plate Method**

Albino mice of either sex (20 - 25g) were grouped as control, standard and test (I-II) containing three numbers. The animals were acclimated to laboratory conditions for one week prior to the experiment. Eddy's hot plate with the temperature of  $55^{\circ} \pm 1^{\circ}\text{C}$  used for the study. Mice were placed on hot plate and the reaction time that is licking of paw or jumping response was recorded in seconds. The cut off time is 15 seconds and the animals not showing any response after 15 seconds were removed from the study. The test compounds were administered intraperitoneally with a dose of 50 mg/kg in 1% polyethylene glycol. Standard group was administered with pentazocine intraperitoneally in a dose of 50mg/kg whereas the control group was given only with 1% v/v polyethylene glycol. The reaction time was measured at the interval of 0, 30, 60, 90 and 180 min. The animal study protocol was approved by IAEC (Reg No: 409/01/CPCSEA). The results are reported in Table No. (Eddy N. B., *et al.*, 1953)



## E. EVALUATION OF ANTIBACTERIAL ACTIVITY

The antibacterial activity can be evaluated by the following techniques.

A) Agar streak dilution method

B) Serial dilution method

C) Agar diffusion method

Cup plate method

Cylinder method

Paper disc method

D) Disc diffusion method

The antibacterial activity of synthesized compounds were screened in the concentration of 25, 50 and 100 µg in dimethyl formamide against gram positive organism *Staphylococcus aureus* and gram negative organism *Escherichia coli* in Muller Hinton agar medium by disc diffusion method. The antimicrobial activity was evaluated by measuring zone of inhibition in mm, the details of the procedure are given below.

### Preparation of Muller Hinton Agar

#### Composition

Beef extract	10.0 g
Casein acid hydrolysis	17.5 g
Starch	1.5 g
Agar	20.0 g
Distilled water	1000 ml

All the ingredients were taken in 1000 ml of distilled water in a conical flask and heated in a steam bath to dissolve. The pH was adjusted to  $7.0 \pm 0.2$  and sterilized in autoclave at 15 lb at  $120^{\circ}\text{C}$  for 15 minutes. The sterile medium was poured into petridish and allowed to solidify.

### Preparation of the Disks

Paper disks of 5 mm diameter and 2 mm thickness were sterilized by autoclaving at  $121^{\circ}\text{C}$  for 15 minutes. Ciprofloxacin ( $10\text{ }\mu\text{g/ml}$ ) was used as standard antibiotic for the comparison of antibacterial activity of the synthesized compounds.

### Organism used


#### Gram positive Organisms

*Staphylococcus aureus* ATCC 6538P

#### Gram negative Organisms

*Escherichia coli* ATCC 2592

### Standard drug



Ciprofloxacin

#### IV. F. EVALUATION OF ANTIFUNGAL ACTIVITY:

The antifungal activity can be evaluated by the following techniques:

Cup and plate method/ Cylinder method

Turbidimetry / Tube assay method

##### Organism used

*Aspergillus niger* ATCC 9029


The antifungal activity of synthesized compounds was screened in the concentration of 25, 50 and 100 µg in dimethyl formamide against *Aspergillus niger*. The antifungal activity was evaluated by measuring zone of inhibition in mm, the details of the procedure are given below.

##### Preparation of Sabouraud's Agar Media

###### Composition

Dextrose	20 g
Peptone	10 g
Purified water	1000 ml
pH	5.4 ± 0.2
Agar	15 g

The media was prepared by dissolving the specified quantities of the dehydrated ingredients (Hi-media) in purified water and was distributed in petridish to a thickness of 3-4 mm. The plates were sterilized by autoclaving at 121 °C for 15 minutes. The sterile medium was poured into petridish and allowed to solidify.

**Standard drug**

Ketoconazole

**Procedure for Antimicrobial activity**

The synthesized compounds were tested for antibacterial and antifungal activity by disc diffusion method at the concentration of 25, 50 and 100  $\mu\text{g/ml}$ . Whatmann filter paper (grade-1) disc of 5 mm diameter and 2 mm width was sterilized by autoclaving for 15 min at 121°C. The sterile agar petridish was seeded with test bacteria and the impregnated discs were placed on the medium with suitable space between the discs. The plates were incubated at  $37^\circ\text{C} \pm 1^\circ\text{C}$  for 18-24 h for bacterial medium and  $25^\circ\text{C} \pm 1^\circ\text{C}$  for 72 h for fungal medium. The inhibition of zones caused by various synthesized compounds and standard drugs ciprofloxacin and ketoconazole on the bacterial and fungal microorganisms were examined.

(*Ind. Pharmacopoeia*, vol -I, 2007).

# RESULTS AND DISCUSSION

## V. RESULTS AND DISCUSSION

The completion of reaction was monitored by TLC using silica gel as stationary phase and chloroform-methanol as mobile phase. The spot in the TLC plate was detected by iodine vapours. The chloroform and petroleum ether mixture was used to purify the compounds. The satisfactory yield was obtained for every reaction. The Physical and spectral data's of the synthesized compounds are given below

### a. Physical data of the compounds

#### i) Physical Data of BOC-Amino acid

The amino group of amino acid is protected by tert-butyl oxy carbonic anhydride (BOC). The BOC amino acids were obtained with good yield. The Physical Data of the BOC-Amino acids are given below.

**Table-2**

BOC-amino acid	Molecular formula	Molecular weight	Physical state	M. P (°C)	Yield (%)
BOC-Leu	C <sub>11</sub> H <sub>21</sub> NO <sub>4</sub>	231.29	White powder	82-85	42.2
BOC - Ala	C <sub>8</sub> H <sub>15</sub> NO <sub>4</sub>	189.21	White powder	110-113	62.0
BOC- Phe	C <sub>14</sub> H <sub>19</sub> NO <sub>4</sub>	265.31	white powder	140-142	72.2
BOC-Gly	C <sub>7</sub> H <sub>13</sub> NO <sub>4</sub>	175.80	White powder	88-90	26.2

## ii) Physical Data of 2-chloromethyl-3-amino acid substituted quinazolin-4-(3H)-ones

The BOC protected amino groups are deprotected by using trifluoro acetic acid in chloroform and the physical data of the compounds are given below

**Table-3**

Compound	Molecular formula	Molecular weight	Physcial State	Yield (%)	R <sub>f</sub> Value
<b>QAL</b>	C <sub>15</sub> H <sub>9</sub> ClN <sub>4</sub> O <sub>2</sub>	322.78	Semisolid	52.12	0.58
<b>QAA</b>	C <sub>12</sub> H <sub>13</sub> ClN <sub>4</sub> O <sub>2</sub>	280.71	Semisolid	70.20	0.64
<b>QAPA</b>	C <sub>18</sub> H <sub>17</sub> ClN <sub>4</sub> O <sub>2</sub>	356.80	Semisolid	85.00	0.68
<b>QAG</b>	C <sub>11</sub> H <sub>11</sub> ClN <sub>4</sub> O <sub>2</sub>	266.68	Semisolid	70.80	0.55

Solvent system for TLC:- Chloroform : Methanol (7:3)

## iii) Physical Data of 2-chloromethyl-3-heterocyclic substituted quinazolin-4(3H)-ones

**Table-4**

Compound	Molecular formula	Molecular weight	Physcial state	Yield (%)	M.P (°C)	R <sub>f</sub> Value
<b>QBEN</b>	C <sub>16</sub> H <sub>9</sub> ClN <sub>4</sub> SO <sub>3</sub>	371.50	Brown colour	70.16	286	0.60
<b>QIBM</b>	C <sub>16</sub> H <sub>11</sub> ClN <sub>4</sub> O	310.00	Brown colour	60.20	194	0.50
<b>QNIBM</b>	C <sub>22</sub> H <sub>15</sub> ClN <sub>5</sub> O <sub>3</sub>	432.83	Brown colour	65.00	274	0.80
<b>QPIN</b>	C <sub>14</sub> H <sub>9</sub> ClN <sub>4</sub> O <sub>3</sub>	316.70	Brown colour	52.00	254	0.70
<b>QBrP</b>	C <sub>14</sub> H <sub>9</sub> BrClN <sub>3</sub> O	350.59	Brown colour	46.00	164	0.64

Solvent system for TLC: - Chloroform: Methanol (7:3)


## b. Spectral analysis

The synthesized compounds were characterized by IR, proton NMR and Mass spectrophotometer and the structures of the compounds are consistent with the spectra.

### IR Spectral Analysis

The IR spectrum was recorded in JASCO FT-IR spectrophotometer. The significant IR values are measured in  $\text{cm}^{-1}$  and the results given in the table.

### Interpretation of IR spectra of QAL



The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-5**

S. No	Wave numbers ( $\text{cm}^{-1}$ )	Functional groups
1	3449	NH Stretching ( $\text{NH}_2$ )
2	2959	Aromatic $-\text{CH}-$ Stretching
3	1649	$\text{C}=\text{O}$ Stretching in amide
4	1515	$\text{C}=\text{N}$ Stretching
5	1368	$\text{C}-\text{N}$ Stretching
6	771	$\text{C}-\text{Cl}$ Stretching







Figure 3: IR Spectra of QAL

### Interpretation of IR spectra of QAA



The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-6**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3435	NH Stretching (NH <sub>2</sub> )
2	2978	Aromatic –CH- Stretching
3	1697	C=O Stretching in amide
4	1585	C=N Stretching
5	1296	C-N Stretching
6	667	C-Cl Stretching





Figure 4: IR Spectra of QAA

**Interpretation of IR spectra of QAPA**

The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-7**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3434	NH Stretching (NH <sub>2</sub> )
2	1524	NH- Stretching
3	1683	C=O Stretching in amide
4	1524	C=N Stretching
5	1299	C-N Stretching
6	700	C-Cl Stretching





Figure 5: IR Spectra of QAPA

### Interpretation of IR spectra of QAG



The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-8**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3431	NH Stretching (NH <sub>2</sub> )
2	1524	NH- Stretching
3	1686	C=O Stretching in amide
4	1585	C=N Stretching
5	1289	C-N Stretching
6	771	C-Cl Stretching

IDEAL ANALYTICAL AND RESEARCH INSTITUTION, PUDUCHERRY





Figure 6: IR Spectra of QAG

**Interpretation of IR spectra of QBEN**

The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-9**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3094	Aromatic-C-H Stretching
2	1647	C=O Stretching in amide
3	1570	C=N Stretching
4	1330	C-N Stretching
5	720	C-Cl Stretching
6	616	C-S bending







Figure 7: IR Spectra of QBEN

**Interpretation of IR spectra of QIBM**

The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-10**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3483	N-H Stretching
2	3132	C-H Aromatic stretching
3	3055	Imidazole
4	1666	C=O Stretching in amide
5	1570	C=N Stretching
6	1566	C=C stretching
7	1313	C-N stretching
8	697	C-Cl stretching

IDEAL ANALYTICAL AND RESEARCH INSTITUTION, PUDUCHERRY.





Figure 8: IR Spectra of QIBM

**Interpretation of IR spectra of QNIBM**

The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-11**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3446	N-H Stretching
2	3102	C-H Aromatic stretching
3	1676	C=O Stretching in cyclic amide
4	1646	C=C Stretching
5	1528	C=N stretching
6	1343	C-N stretching
7	758	C-Cl stretching

IDEAL ANALYTICAL AND RESEARCH INSTITUTION, PUDUCHERRY





Figure 9: IR Spectra of QNIBM

**Interpretation of IR spectra of QPIN**

The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-12**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3102	C-H Aromatic stretching
2	1676	C=O Stretching in cyclic amide
3	1606	C=C Stretching
4	1588	C=N stretching
5	1243	C-N stretching
6	1530	NO <sub>2</sub> Asymmetrical stretching
7	752	C-Cl stretching

IDEAL ANALYTICAL AND RESEARCH INSTITUTION, PUDUCHERRY.





Figure 10: IR Spectra of QPIN

**Interpretation of IR spectra of QBrP**

The significant wave numbers of the compound and its relevant functional groups are given below:

**Table-13**

S. No	Wave numbers (cm <sup>-1</sup> )	Functional groups
1	3242	C-H Aromatic stretching
2	1656	C=O Stretching in amide
3	1602	C=C Stretching
4	1586	C=N stretching
5	752	C-Cl stretching
6	650	C-Br bending



IDEAL ANALYTICAL AND RESEARCH INSTITUTION, PUDUCHERRY





Figure 11: IR Spectra of QBrP

## ii) NMR Spectral Analysis

The proton NMR values were recorded in the BRUKER 400 MHz NMR spectrophotometer

### Interpretation of $^1\text{H}$ NMR spectra of QAL



The  $\delta$  values with reference to the nature of protons are given below.

**Table-14**

S. No	Values in ppm	Nature of protons
1	8.1	Protons of aromatic ring in quinazolone
2	7.2	Protons in Amidic nitrogen
3	5.3	Protons in chloromethyl
4	1.5	Protons of primary amine
5	0.9	Proton of isopropyl group




Figure 12: NMR Spectra of QAL




Figure 13: NMR Spectra of QAL




Figure 14: NMR Spectra of QAL

Q-AL




Figure 15: NMR Spectra of QAL





Figure 16: NMR Spectra of QAL

**Interpretation of  $^1\text{H}$  NMR spectra of QAA**

The  $\delta$  values with reference to the nature of protons are given below.

**Table-15**

S. No	Values in ppm	Nature of protons
1	8.1	Protons of aromatic ring in quinazalone
2	7.3	Protons of Amidic nitrogen
3	5.4	Protons in chloromethethyl
4	3.6	Alkyl proton of methine group
5	1.9	Protons of primary amine
6	1.3	Alkyl proton of methyl group
7	1.0	Proton of isopropyl group






Figure 17: NMR Spectra of QAA

Q-AA





Figure 18: NMR Spectra of QAA

Q-AA



BRUKER  
 AVANCE II 400 NMR  
 Spectrometer  
 SAIF  
 Panjab University  
 Chandigarh

Current Data Parameters  
 NAME Dec19-2011  
 EXPNO 280  
 PROCNO 1


F2 - Acquisition Parameters  
 Date\_ 20111219  
 Time 16.51  
 INSTRUM spect  
 PROBHD 5 mm PABBO BB-  
 PULPROG zg30  
 TD 65536  
 SOLVENT CDCl3  
 NS 8  
 DS 2  
 SWH 12019.230 Hz  
 FIDRES 0.183399 Hz  
 AQ 2.7263477 sec  
 RG 40.3  
 DW 41.600 usec  
 DE 6.00 usec  
 TE 294.2 K  
 D1 1.00000000 sec  
 TDO 1

===== CHANNEL f1 =====  
 NUC1 1H  
 P1 10.90 usec  
 PL1 -3.00 dB  
 SFO1 400.1324710 MHz

F2 - Processing parameters  
 SI 32768  
 SF 400.1299789 MHz  
 WDW EM  
 SSB 0  
 LB 0.30 Hz  
 GB 0  
 PC 1.00

avtar\_saifpu@yahoo.co.in


Figure 19: NMR Spectra of QAA

**Interpretation of  $^1\text{H}$  NMR spectra of QAPA**

The  $\delta$  values with reference to the nature of protons are given below.

**Table-16**

S. No	Values in ppm	Nature of protons
1	7.31	Protons of aromatic ring in quinazolone
2	7.18	Protons of free phenyl group in quinazolone
3	6.8	Protons of amidic nitrogen
4	5.1	Protons in chloromethethyl
5	1.7	Protons of primary amine
6	1.3	Alkyl proton of methyl group



### Figure 20: NMR Spectra of QAPA




Figure 21: NMR Spectra of QAPA





Figure 22: NMR Spectra of QAPA

Q-APA



BRUKER  
 AVANCE II 400 NMR  
 Spectrometer  
 SAIF  
 Panjab University  
 Chandigarh

Current Data Parameters  
 NAME Dec19-2011  
 EXPNO 270  
 PROCNO 1

F2 - Acquisition Parameters  
 Date\_ 20111219  
 Time 16.46  
 INSTRUM spect  
 PROBHD 5 mm PABBO BB-  
 PULPROG zg30  
 TD 65536  
 SOLVENT CDCl3  
 NS 8  
 DS 2  
 SWH 12019.230 Hz  
 FIDRES 0.183399 Hz  
 AQ 2.7263477 sec  
 RG 57  
 DW 41.600 usec  
 DE 6.00 usec  
 TE 294.2 K  
 D1 1.00000000 sec  
 TD0 1

===== CHANNEL f1 =====  
 NUC1 1H  
 P1 10.90 usec  
 PL1 -3.00 dB  
 SFO1 400.1324710 MHz

F2 - Processing parameters  
 SI 32768  
 SF 400.1300024 MHz  
 WDW EM  
 SSB 0  
 LB 0.30 Hz  
 GB 0  
 PC 1.00

avtar\_saifpu@yahoo.co.in

Figure 23: NMR Spectra of QAPA







Figure 24: NMR Spectra of QAPA

**Interpretation of  $^1\text{H}$  NMR spectra of QAG**

The  $\delta$  values with reference to the nature of protons are given below.

**Table-17**

S. No	Values in ppm	Nature of protons
1	8.02	Protons of amidic nitrogen
2	7.4	Protons of aromatic ring in quinazolone
3	4.1-4.4	Protons in chloromethyl
4	3.8	proton of methyl group
5	1.5	Protons of primary amine




Figure 25: NMR Spectra of QAG

Q-AG




Figure 26: NMR Spectra of QAG





Figure 27: NMR Spectra of QAG

**Interpretation of  $^1\text{H}$  NMR spectra of QBEN**

The  $\delta$  values with reference to the nature of protons are given below.

**Table-18**

S. No	Values in ppm	Nature of protons
1	8.0 – 8.1	Protons of aromatic ring in benzthiazole
2	7.4 – 7.7	Protons of aromatic ring in quinazolone
3	4.0	Protons in chloromethyl




Figure 28: NMR Spectra of QBEN






Figure 29: NMR Spectra of QUBEN





**Figure 30: NMR Spectra of QBEN**

**Interpretation of  $^1\text{H}$  NMR spectra of QIBM**

The  $\delta$  values with reference to the nature of protons are given below.

**Table-19**

S. No	Values in ppm	Nature of protons
1	7.8	Protons of aromatic ring in quinazolone
2	7.1	Protons of aromatic ring in Benzimidazole
3	5.2	Protons in chloromethethyl
4	5.0	Protons of secondary amine




Figure 31: NMR Spectra of QIBM





Figure 32: NMR Spectra of QIBM

### Interpretation of $^1\text{H}$ NMR spectra of QNIBM



The  $\delta$  values with reference to the nature of protons are given below.

**Table-20**

S. No	Values in ppm	Nature of protons
1	8.1	Protons of aromatic ring in quinazolone
2	7.3–7.8	Protons in phenyl group
3	7.0	Protons of aromatic ring in Benzimidazole
4	4.1-4.3	Protons in chloromethethyl
5	5.0	Protons of secondary amine




Figure 33: NMR Spectra of QNIBM




Figure 34: NMR Spectra of QNIBM






Figure 35: NMR Spectra of QNIBM





### Figure 36: NMR Spectra of QNIBM

**Interpretation of  $^1\text{H}$  NMR spectra of QPIN**

The  $\delta$  values with reference to the nature of protons are given below.

**Table-21**

S. No	Values in ppm	Nature of protons
1	8.5	Protons of aromatic ring in pyridine
2	7.7-8.1	Protons of aromatic ring in quinazolone
3	4.1-4.4	Protons in chloromethyl




Figure 37: NMR Spectra of QPIN





Figure 38: NMR Spectra of QPIN

**Interpretation of  $^1\text{H}$  NMR spectra of QBrP**

The  $\delta$  values with reference to the nature of protons are given below.

**Table-22**

S. No	Values in ppm	Nature of protons
1	8.6	Protons of aromatic ring in pyridine
2	8.1	Protons of aromatic ring in quinazolone
3	4.1-4.4	Protons in chloromethyl




Figure 39: NMR Spectra of QPIN




Figure 40: NMR Spectra of QBrP




Figure 41: NMR Spectra of QBrP







Figure 42: NMR Spectra of QBrP

### iii) Mass Spectral Analysis

Mass spectrum of the sample was recorded in JEOL GCmate by Electron impact method as ionization mode.

#### Interpretation of Mass spectra of QAL




Molecular Weight: 322.78

Molecular Ion Peak: 322.78


The possible fragments of the molecule with the relevant to its m/z values are:

**Table-23**

S. No	m/z	Fragments
1	309.52	
2	281.70	
3	252.61	

**Figure 43: Mass Spectra of QAL**

## Interpretation of Mass spectra of QAA




Molecular Weight: 280.71

Molecular Ion Peak: 280.71


The possible fragments of the molecule with the relevant to its m/z values are:

Table-24

S. No	m/z	Fragments
1	251.62	<chem>CC(=O)Nc1nc(CCl)c2ccccc12=O</chem>
2	218.62	<chem>Nc1nc(CCl)c2ccccc12=O</chem>
3	184.71	<chem>Cc1nc(N)c2ccccc12=O</chem>
4	148	<chem>Nc1nc2ccccc2c(=O)[nH]1</chem>

**Figure 44: Mass Spectra of QAA**

## Interpretation of Mass spectra of QAPA




Molecular Weight: 356.80

Molecular Ion Peak: 356.80


The possible fragments of the molecule with the relevant to its m/z values are:

Table-25

S. No	m/z	Fragments
1	296.73	
2	282.72	
3	253.16	
4	208.69	
5	122.63	

**Figure 45: Mass Spectra of QAPA**

## Interpretation of Mass spectra of QAG



Molecular Weight: 266.34


Molecular Ion Peak: 266.68

The possible fragments of the molecule with the relevant to its m/z values are:


Table-26

S. No	m/z	Fragments
1	253.56	<chem>ClCC1=NC2=CC=CC=C2C(=O)N1NCCN</chem>
2	236.55	<chem>CCCN1=NC2=CC=CC=C2C(=O)N1C(Cl)C</chem>
3	208.62	<chem>NC1=NC2=CC=CC=C2C(=O)N1C(Cl)C</chem>
4	194.61	<chem>ClCC1=NC2=CC=CC=C2C(=O)N1</chem>



**Figure 46: Mass Spectra of QAG**

## Interpretation of Mass spectra of QBEN



Molecular Weight: 325.87

Molecular Ion Peak: 325.41

The possible fragments of the molecule with the relevant to its m/z values are:

Table-27

S. No	m/z	Fragments
1	275.3	
2	141.4	
3	133.5	
4	119.4	





Figure 47: Mass Spectra of QBEN

### Interpretation of Mass spectra of QIBM




Molecular Weight: 274.27

Molecular Ion Peak: 274.36


The possible fragments of the molecule with the relevant to its m/z values are:

**Table-28**

S. No	m/z	Fragments
1	261	
2	236	
3	145	
4	117	

**Figure 48: Mass Spectra of QIBM**

## Interpretation of Mass spectra of QNIBM



Molecular Weight: 432.31

Molecular Ion Peak: 432.83

The possible fragments of the molecule with the relevant to its m/z values are:

Table-29

S. No	m/z	Fragments
1	389.5	
2	350.78	
3	312.74	
4	284.72	
5	210.66	





Figure 49: Mass Spectra of QNIBM

## Interpretation of Mass spectra of QPIN



Molecular Weight: 316.45

Molecular Ion Peak: 316.59

The possible fragments of the molecule with the relevant to its m/z values are:

Table-30

S. No	m/z	Fragments
1	268.22	<chem>O=C1Nc2ccccc2N1c3ccc([N+](=O)[O-])cc3</chem>
2	257.24	<chem>C=CC1=NC2=CC=CC=C2C(=O)N1c3ccc([N+](=O)[O-])cc3</chem>
3	218.25	<chem>C=CC(NCc1ccccc1)c2ccc([N+](=O)[O-])cc2</chem>
4	173.25	<chem>C=CC(NCc1ccccc1)</chem>
5	107.15	<chem>NCCc1ccccc1</chem>







Figure 50: Mass Spectra of QPIN

## Interpretation of Mass spectra of QBrP



Molecular Weight: 350.57

Molecular Ion Peak: 350.59

The possible fragments of the molecule with the relevant to its m/z values are:

Table-31

S. No	m/z	Fragments
1	318	
2	306	
3	296	
4	253	
5	236	




Figure 51: Mass Spectra of QBrP

### Evaluation of analgesic activity

The analgesic activity of the synthesized compounds was evaluated by Eddy's hot plate method in mice with reference to the standard drug pentazocine.

**Table-32. Results of Analgesic activity**

Compound	Paw licking or jumping response (in sec)				
	0 min	30 mins	60 mins	90 mins	180 mins
Control	7.02 ±0.58	8.06 ±0.58	7.08 ±0.58	7.01 ±0.58	6.67 ±0.67
Standard (50mg/kg)	10.04±1.15	11.03 ±0.58	10.21 ±0.58	9.67 ±0.33	8.27 ±0.88
QAPA (50mg/kg)	7.02 ±0.78	9.76±0.32**	8.88±0.58**	8.78 ±0.78*	6.08±0.98
QIBM (50mg/kg)	8.06 ±0.58	9.33 ±0.33	9.73 ±0.33*	7.02±0.58	6.33 ±0.31

\*P<0.05, \*\*P<0.01. \*\*\*P<0.001 compare to 0 h response. One-way ANOVA followed by Bonferroni test.



## Screening of antimicrobial activity

### i) Antibacterial Activity



The antibacterial activity of synthesized compounds were screened against both gram positive (*S. aureus*) and gram negative (*E. coli*) organism by agar disc diffusion method using ciprofloxacin as standard drug.

**Table-33. Results of Antibacterial Activity**

Compounds	Zone of Inhibition (in mm)					
	<i>S. aureus</i>			<i>E. coli</i>		
	25 (µg/ml)	50 (µg/ml)	100 (µg/ml)	25 (µg/ml)	50 (µg/ml)	100 (µg/ml)
<b>QAL</b>	22	26	31	11	14	20
<b>QAA</b>	19	22	26	19	22	24
<b>QAPA</b>	15	19	23	21	24	29
<b>QAG</b>	14	19	22	18	21	26
<b>QBEN</b>	15	18	27	20	23	29
<b>QIBM</b>	19	21	27	16	21	27
<b>QNIBM</b>	16	19	29	17	22	28
<b>QPIN</b>	22	26	28	20	22	28
<b>QBrP</b>	20	22	28	15	18	24
<b>Ciprofloxacin 10(µg/ml)</b>	39			38		



**Figure 52: Antimicrobial activity of synthesized compounds against *S. aureus***



**Figure 53: Antimicrobial activity of synthesized compounds against *E. coli***



## ii) Antifungal Activity

The antifungal activities of synthesized compounds were screened against *A.niger* by agar disc diffusion method using ketoconazole as standard drug.

**Table-34. Results of Antifungal activity**

Compounds	Zone of Inhibition (in mm)		
	<i>Aspergillus niger</i>		
	25 (µg/ml)	50 (µg/ml)	100 (µg/ml)
<b>QAL</b>	22	26	31
<b>QAA</b>	19	22	26
<b>QAPA</b>	15	19	23
<b>QAG</b>	14	19	22
<b>QBEN</b>	15	19	23
<b>QIBM</b>	13	16	18
<b>QNIBM</b>	21	24	29
<b>QPIN</b>	22	26	28
<b>QBrP</b>	15	13	24
<b>Ketoconazole 10(µg/ml)</b>	39		





**Figure 54: Antimicrobial activity of synthesized compounds against *A.niger***

## DISCUSSION

### **Analgesic activity:**

The synthesized compounds were tested for their Analgesic activity by Eddy's hot plate method where the source of pain is heat and the results of the compounds were compared with the standard drug pentazocine. The activity was observed up to 180 minutes.

The standard drug pentazocine showed increase in reaction time to the stimulus with maximum response at 60<sup>th</sup> minute. Also at 60<sup>th</sup> minute, both QAPA and QIBM were found to be significant compared to control. The activity was started to abolish gradually after 60<sup>th</sup> min. One-way ANOVA followed by Bonferroni test was applied to the results to find the results are statistically significant or not in the used population.

### **Antimicrobial activity**

The antimicrobial activity was screened disc diffusion method were Muller Hinton Agar media used for bacteria and Sabouraud's agar media used for fungal growth.

### **Antibacterial activity**

Among the compounds used for antibacterial activity, QPIN, QNIBM, QBEN could show significant activity against both (*S. aureus*) and (*E. coli*) organisms. The QAL and QAA were somewhat selective for the inhibition of gram positive organism (*S. aureus*). The rest of the compounds could show only moderate activity at the optimal concentration of 100µg/ml when compared to 10µg/ml of ciprofloxacin as standard.

### **Antifungal activity**

Among the compounds QAL, QNIBM and QPIN shown good antifungal activity against *A. niger* and rest of the compounds could show only moderate activity at the maximum concentration of 100µg/ml when compared to 10µg/ml of ketoconazole as standard.

# SUMMARY

## VI. SUMMARY

Totally nine compounds of newly modified quinazolin-4 (3H)-ones were synthesized. Four derivatives of quinazolones were synthesized by incorporating the amino acids of Leucine, alanine, phenylalanine and glycine at third position of the ring. Anthranilic acid was treated with chloroacetyl chloride and obtained N-chloro acetyl anthranilic acid. To synthesize amino acid incorporated quinazolones, the N-chloroacetyl anthranilic acid was treated with hydrazine hydrate and derived 3-amino-2-chloromethyl quinazolin-4(3H)-one. Then this intermediate was coupled with protected amino acids, using EDC as coupling agent. Finally the BOC was deprotected to have 2-chloromethyl-3-amino acid substituted quinazolin-4(3H)-ones.

Five derivatives of quinazolones were synthesized by incorporating the heterocyclic moieties of substituted Indole, Benzthiazole and Pyridine at third position of the ring. Anthranilic acid was treated with chloroacetyl chloride and obtained N-chloro acetyl anthranilic acid. To synthesize heterocyclic incorporated quinazolones, N-Chloroacetyl anthranilic acid was refluxed separately with 2-amino benzthiazole, 2- amino benzimidazole, 2-amino 5-nitro phenyl benzimidazole, 2-amino 5-nitro pyridine and 2-amino 5-bromo pyridine and resulting 2-chloromethyl-3-heterocycle substituted quinazolin-4(3H)-one.

The completion of the reaction was checked by TLC. The chloroform and petroleum ether mixture was used for the purification of the amino acid derivatives of quinazolinones and the heterocyclic derivatives of quinazolinones was washed thoroughly with boiling water and recrystallized from acetone: ethanol mixture (1:1).

The structures of the synthesized compounds were confirmed with IR, <sup>1</sup>H NMR and Mass spectroscopy.

The synthesized compounds were tested for their analgesic activity by Eddy's hot plate method where the source of pain is heat and the results of the compounds were compared with the standard drug pentazocine. One-way ANOVA followed by Bonferroni test was applied to the results to find the results are statistically or not in the used population.

The synthesized compounds were screened for their antimicrobial activity for the gram positive, gram negative and fungal organism by disc diffusion method and the inhibition of microorganism were compared with the standard drugs ciprofloxacin and ketoconazole respectively.

The compounds QAPA, QIBM shown moderate analgesic activity.

The compounds QAL, QAA, QBEN, QNIBM and QPIN were shown significant activity against the organism used for the study.

# CONCLUSION

## VII. CONCLUSION

The two different set of quinazolones containing amino acids and heterocycles separately were synthesized and evaluated. The analgesic and antimicrobial activities are not very significant for one set of compounds than the other. Though all the compounds shown biological action it is better to incorporate the dipeptides and other heterocycles.

For the amino acid series the rationale cannot be drawn for the different amino acids used, since the aliphatic amino acid leucine and alanine activity is equal to aromatic amino acid phenyl alanine. There is no distinct difference in the biological action upon increasing the hydrophobicity through increasing number of carbons. May be the di or tripeptide incorporation at the 3<sup>rd</sup> position can give an idea about whether the amino acid/peptides incorporation in the quinazolones may overtake the heterocycles or not.

For heterocyclic incorporate series, among the Benzimidazole and Benzthiazole and pyridine used, the nitro group substituted compounds are active than others. May be the electron withdrawing group are required to enhance the biological action of the quinazolones.



# BIBLIOGRAPHY

## VIII.BIBLIOGRAPHY

- ❖ Ashis Kumar Nanda., Subarna Ganguly. and Ranadhir Chakraborty. Antibacterial activity of some 3-(Arylidene amino)-2-phenyl quinazolin-4(3H)-ones synthesis and Preliminary QSAR Studies, *Molecules*, **2007**, 12, 2413-2426.
- ❖ Ashok Kumar., Shalabh Sharma., Archana., Kiran Bajaj., Shipra Sharma., Hemant Panwar., Tripti Singh. and Srivastava V. K. Some new 2, 3, 6-trisubstituted quinazolinones as potent antiinflammatory, analgesic and COX-II inhibitors, *Bioorganic & Medicinal Chemistry*, **2003**, 11, 5293–5299.
- ❖ Avinash Patil., Swastika Ganguly. and Sanjay Surana. Synthesis of 2-[(5-substituted-1H-benzimidazol-2-yl sulfinyl) methyl]-3-substituted Phenyl Quinazolin-4(3H)-one derivatives, *Asian Journal of Chemistry*, **2009**, 21(2), 1490-1494.
- ❖ Beckett A. H. and Stenlake J. B. Practical Pharmaceutical Chemistry, 4<sup>th</sup> edn, part-II, S.K Jain for CBS Publishers, New Delhi. **2007**, 415-456.
- ❖ Beena K. P. and Akelesh T. Synthesis and antibacterial activity of quinazolinone derivatives, *International Journal of Pharmacy and Pharmaceutical Sciences*, **2010**, 2(3), 166-168.
- ❖ Bodanszky and Sodanszky M. *Practice of Peptide Synthesis*, Springer verlog, Newyork. **1984**, 78.
- ❖ Chatrasal Singh Rajput., Ashok Kumar., Sudhir Kumar Bhati. and Jai Singh. Synthesis and antiinflammatory activity of 2-{5'-(4-Pyridinyl)-1', 2', 3' -

- oxadiazol-2-yl-thiomethyl]-3-substituted-aryl-6-substituted-quinazolin-4-one, *Asian Journal of Chemistry*, **2008**, 20 (8), 6246-6252.
- ❖ Deb A. C. Fundamentals of Biochemistry, 9<sup>th</sup> edn, New Central Book Agency (p) Ltd., Kolkata. **2008**, 82-91.
  - ❖ Gowramma B., Jubie S., Kalirajan R. and Suresh B. Synthesis of quinazolinone derivatives with nitrogen mustard as possible anticancer agents, *Asian Journal of Chemistry*, **2009**, 21 (7), 5235-5240.
  - ❖ Gupta S. K, *Drug Screening Methods*, Japee Brother Publisher, **2004**, 151.
  - ❖ Indian Pharmacopoeia, Central Indian Pharmacopoeia Laboratory Govt. of India, Ministry of Health & Family Welfare, New Delhi, **2007**, (1), 45-57.
  - ❖ Jag Mohan, Organic Spectroscopy, Principles and Applications, 2<sup>nd</sup> edn, Narosa publishing home, New Delhi. **2005**, 76-95.
  - ❖ Jayshari S. Pattan., Shashikant R. Pattan., Nachiket S Dighe., Hariprasad C. K., Nirmal S.A. and Hiremath S. N. Synthesis and evaluation of some new quinazolone derivatives for their antimicrobial activity, *International Journal of Pharmaceutical research and Development*, **2009**, 1 (9), 1-8.
  - ❖ Jessy E. M., Thirugnanasambanthan A., Alex J., Sridevi H. and Srinivasan K. K. Synthesis and biological evaluation of some novel quinazolones, *Indian Journal of Pharmaceutical Sciences*, **2007**, 69 (3), 476-478.
  - ❖ Jitender Singh., Parul Grover. and Dharam Pal Pathak. Synthesis and Comparative QSAR study of some novel benzimidazole derivatives, *Acta pharmaceutica Scientia*, **2010**, 52, 510-521.

- ❖ Kale A. U., Kardile D. P., Kalyane N. V., Patel M. R. and Patel H. Synthesis and antimicrobial activity of some 2,3-disubstituted quinazoline-4(3H)-ones, *International Journal of Pharmaceutical and Applied Sciences*, **2010**, 1 (1), 85-90.
- ❖ Kulkarni S. K, Hand Book of Experimental Pharmacology, 3<sup>rd</sup> edn, Vallabh Prakashan, New Delhi. **2007**, 128-130.
- ❖ Manish Srivastav., Salahuddin M. D. and Shantakumar S. M. Synthesis and antiinflammatory activity some novel 3-(6-Substituted-1, 3-benzothiazol-2-yl)-2-[(4-substituted phenyl) amino] methyl quinazolin-4(3H)-ones, *E- Journal of Chemistry*, **2009**, 6 (4), 1055-1062.
- ❖ Mariappan G., Saha B. P., Satyajit Dutta., Abhishek Majumder. and Sudip Saha. Synthesis and biological evaluation of quinazolone Fused Schiff Bases, *Indian Journal of Pharmaceutical Education and Research*, **2011**, 45(1), 78-82.
- ❖ Narendra Babu A. and Rama Rao Nadendla. Synthesis of some new quinazolinone formazans as antiinflammatory and anthelmintic agents, *Journal of Pharmacy Research*, **2011**, 4 (4), 983-985.
- ❖ Palle Acharyulu V. R., Dubey P. K., Prasada reddy P. V. V. and Thatipally Suresh. Synthesis of 2-(4-substituted sulfonyl piperazin-1-ylmethyl-3-aryl- quinazolin-4(3H)-one, *Indian Journal of Chemistry*, **2010**, 48B, 923-928.
- ❖ Pandey V. K., Pathak L. P. and Mishra S. K. Synthesis a new series of isoquinoliny quinazolones, antiviral and antifungal activities, *Indian Journal of Chemistry*, **2005**, 43B, 1940-1943.

- ❖ Patel J. A., Mistry B. D. and Desai K. R. Synthesis and antimicrobial activity of newer quinazolinones, *E-Journal of Chemistry*, **2006**, 3 (11), 97-102.
- ❖ Ponnilaravasan Ilangoan., Swastika ganguly. and Vijay pandi. Design and synthesis of novel quinazolinone derivatives as broad spectrum anticonvulsant and antimicrobial agent, *Journal of Pharmacy Research*, **2010**, 3 (4), 703-706.
- ❖ Raghavendra N. M., Thampi P. P. and Gurubasavarajaswamy P. M. Synthesis and antimicrobial activity of some novel substituted piperazinyl-quinazolin-3(4*H*)-ones, *E-Journal of Chemistry*, 5( 1), **2008**, 23-33.
- ❖ Rajveer C. H., Stephen Rathinaraj B., Kumaraswamy D., Sudharshini S., Swarnalatha C. H. and Preetha. S. Panicker. Synthesis and biological evaluation of some novel oxo-quinazoline derivatives for their antibacterial activity, *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, **2010**, 1 (2), 366-371.
- ❖ Rang H. P., Dale M. M., Ritter J. M., Flower R. J., Rang and Dale's Pharmacology, 6<sup>th</sup> edn, Churchill Livingstone Elsevier, China. **2008**, 588-609.
- ❖ Robert M. Silverstein, Francisx. Webster, Spectrometric Identification of organic compounds , 6<sup>th</sup> edn, John Wiley & Sons, Inc. **1998**, 15-70.
- ❖ Satyanarayana U, Biochemistry, Arunabhasen Publication, Kolkata. **2005**, 49-49.
- ❖ Sharma Y. R, Elementary Organic Spectroscopy, 4th edn, S. Chand and company Limited, New Delhi. **2007**, 210-233.

- ❖ Shashikant R. Pattan., Krishna Reddy V. V., Manvi F. V., Desai B. G. and Bhat A. R. Synthesis of N-3(4-(4-chlorophenyl thiazole-2-yl)-(2- (amino)methyl)-quinazolin-4(3H)-one and their derivatives for antitubercular activity, *Indian Journal of Chemistry*, **2006**, 45B, 1778-1781.
- ❖ Sowjanya C., Rajitha G., RamaBharathi V. and Kalpana Devi G. Synthesis and evaluation of some novel 3-[5-phenyl-1,3,4-oxadiazol-2-yl]-2-(substituted styryl)-quinazoline-4(3H)-ones for antibacterial activity, *Journal of Chemical and Pharmaceutical Research*, **2011**, 3 (6), 212-216
- ❖ Sreena K., Ratheesh R., Rachana M., Poornima M. and Shyni C. Synthesis and anthelmintic activity of benzimidazole derivatives, *Hygeia*, **2009**, 1 (1), 21-22.
- ❖ Sridhar V., Venugopal G., Venkateshwara Rao., Pranay kumar K., Murthy U. and RaghuRam Rao A. Synthesis of novel 2, 4, 6 trisubstituted quinazoline derivatives showing antibacterial and cytotoxic activities against THP-1, HL-60 and A375 cell line, *Indian Journal of Chemistry*, **2009**, 48B, 840-847.
- ❖ Srinivasa Reddy P., Venugopala K. N., Gopal Krishna Rao. and sanjay Pai. Synthesis of some substituted quinazolinones as analgesic and antiinflammatory agents, *Indian Journal of Heterocyclic Chemistry*, **2007**, 16, 243-246.
- ❖ Suthakaran R., Kavimani S., Venkaiaiah P. and Suganthi K. Synthesis of 3-(2-(4)-4-substitutedbenzylidene-4,5-dihydro-5-oxo-2-phenylimidazol-1-yl)ethyl)-6,8-un/dibromosubstituted-2-substituted quinazoline-(3H)-one, *Rasayan Journal of Chemistry*, **2008**, 1 (1), 22-29.

- ❖ Tripti Singh., Shalabh sharma., Virendra Kishore., Srivastava. and Ashok Kumar. Synthesis insecticidal and antimicrobial activities of some heterocyclic derivatives of quinazolinone, *Indian Journal of Chemistry*, **2006**, 45B, 2558-2565.
- ❖ Vijai Anand P. R., Suresh Kumar K., Sivakumar R., Sam Solomon W. D. and Jayaveera K. N. Synthesis of quinazoline derivatives and their Biological activities, *Asian Journal of Chemistry*, **2009**, 21 (9), 6656-6660.
- ❖ Vogel I, Elementary Practical Organic Chemistry Part I: small Scale preparations, 2<sup>nd</sup> edn, S. K Jain CBS Publishers, New Delhi. **2004**, 249-250.
- ❖ William Kemp, Organic Spectroscopy, 3<sup>rd</sup> edn, Palgrave Publishers, New York. **2007**, 171-176.

# ANNEXURE



## ANNEXURE I

The animal study protocol was approved by IAEC (Reg No: 409/01/CPCSEA).

**CERTIFICATE**

This is certify that the project title SCREENING OF ANALGESIC  
ACTIVITY FOR 2,3-DISUBSTITUTED QUINAZOLONE  
DERIVATIVES.....has been approved by the  
IAEC.

S. SHUBA  
Name of Chairman/member Secretary IAEC

Dr P. Balakrishna murthy  
Name of CPCSEA nominee

Signature with date

S. Shuba 20/7/2011  
Chairman/Member Secretary of IAEC

P. Balakrishna murthy  
CPCSEA nominee

(Kindly make sure that minutes of the meeting duly signed by all the participants are maintained by office)

**[ATTESTED]**

Dr. T. Vetrivelan, M.Pharm., Ph.D.,  
Professor and Principal,  
Adhiparasakthi College of Pharmacy,  
Mulluruthur-603 019,  
Tamilnadu, India.

## ANNEXURE II

Available online [www.jocpr.com](http://www.jocpr.com)

Journal of Chemical and Pharmaceutical Research, 2012, 4(2):1147-1150



Research Article

ISSN : 0975-7384  
CODEN(USA) : JCPRC5**Synthesis of 2, 3-disubstituted quinazolone derivatives for analgesic and antimicrobial activities**

A. Thirugnanasambanthan\* and S. Sankarnarayanan

Adhiparasakthi College of Pharmacy, Melmaruvathur

**ABSTRACT**

*In view of the effective range of biological activities exhibited by quinazolinones, a series of 2, 3-disubstituted quinazolin-4(3H)-ones have been synthesized and evaluated for analgesic, antibacterial and antifungal activities. Five derivatives of quinazolinones were synthesized by incorporating the heterocyclic moieties of substituted Indole, Benzothiazole and Pyridine at third position of the ring. The structures of the compounds were confirmed on the basis of IR, <sup>1</sup>H NMR and Mass spectroscopy. The analgesic and antimicrobial activity of the compounds was screened by Eddy's hot plate method and Disc diffusion method respectively. All the five compounds could show analgesic and antimicrobial activities. The compound II showed better analgesic activity than other compounds. Significant activity was shown by compound IV against gram positive organisms and by compound I against gram negative organisms. The best antifungal activity was shown by compound IV.*

**Keywords:** Quinazolinones, Analgesic, Antibacterial and Antifungal activity.

**INTRODUCTION**

Quinazolin-4(3H)-one is a versatile heterocyclic moiety for the medicinal chemists for molecular modification due to its wide spectrum of biological activities. They have established a variety of biological activities like analgesic [1], anti-inflammatory [1], antihypertensive [2], antihistaminic, anticancer [3-4], sedative- hypnotic and antimicrobial activities [5-6]. The second and third positions are the most favourable positions for making substitutions on the quinazolinone ring and many research works have evidenced this concept [7]. This prompted us to synthesize a new series of quinazolinone derivatives by incorporating the heterocyclic rings at the third position. The 2-chloro methyl-3-heterocycle substituted quinazolinones were synthesized by cyclizing the N-chloro acetyl anthranilic acid with amino substituted indole, benzthiazole and pyridine. The N-chloro acetyl anthranilic acid was prepared by acetylating the anthranilic acid. The structures of all the synthesized compounds were confirmed by IR, <sup>1</sup>H NMR and mass spectroscopy. All the compounds have been screened for their analgesic, antibacterial and antifungal activities.

**EXPERIMENTAL SECTION**

The melting points were determined in open capillary tubes in a Hicon melting point apparatus. Infra-red spectrum was recorded in Jasco FT/IR 410 spectrometer by KBr pellet method and the <sup>1</sup>H NMR spectra was taken on a Bruker Avance II (400 MHz) NMR spectrometer using dimethylsulfoxide as solvent and chemical shifts are expressed in parts per million relative to tetra methyl silane as an internal standard. Mass spectra was recorded on shimadzu 2010A LC-MS. The completion of the reactions was checked by thin layer chromatography on silica gel G coated plates using chloroform: methanol (7:3) solvent system and the spots were detected by Iodine vapours. The animal study protocol was approved by Institutional Animal Ethical Committee. (Reg No: 409/01/CPCSEA).